

Patient Guide



Endoselect Salivary #4119*

Endoselect Salivary NY #4120

*Not Available in New York

BEFORE YOU BEGIN

Activate This Test

Visit gdx.net/activate and enter the number found on the included activation label card.



STEP 1

Plan Your Collection

Use a calendar to plan your specimen collection. Ship Monday thru Friday and avoid US holidays which may cause delays.

Prior to Testing

Certain medications may influence results, however there are no known interfering medications. Do not change use of medications unless instructed to do so by your healthcare provider.

If you have difficulty producing enough saliva for the tube, press the tip of your tongue against your teeth. Yawning, imagining sour or flavorful foods, or smelling flavorful foods can also generate saliva.

One Hour Before Testing

One hour prior to collection do not eat, brush or floss teeth, use mouthwash, lip balm, chewing gum or any tobacco products; you may drink **ONLY** water during this time.

Continued on next page

STEP 1 (Continued from previous page)

Plan Your Collection

Premenopausal Women

If the goal is to evaluate ovulatory function, collect specimens between days 19 and 25 of the menstrual cycle.

Peri/Menopausal Women and Males

Collect specimens on any day.

Patients on Hormone Therapies

Do not take hormones immediately prior to the morning specimen collection. Wait to take the dose until after the specimen has been collected unless otherwise instructed by your healthcare provider.

As a general guideline, many clinicians opt to test patients approximately 8–12 hours following a dose. There are many variables that affect hormone levels with respect to timing of collection including dosing regimen, formulation pharmacokinetics, route of administration, site of administration, length of regimen, patient metabolic individuality, specimen type, etc.

Synthetic, non-bioidentical hormones are not detected by the assays and testing is not recommended for these patients due to questionable clinical utility.

STEP 2

Specimen Collection

1. **Review** instructions and test prep information at gdx.net/activate.
2. Write your **date of birth (DOB)**, **collection start time** and **collection stop time** and the **date of collection** on the labels provided. Attach completed label to the saliva tube.



Collect Saliva





1. Refer to the chart at right for when to begin saliva collection.
2. When ready to collect the saliva, **rinse your mouth with drinking water**. Spit out the water completely, and **do not cough or clear your throat** into the collection tube. **Discard** the saliva **if there is visible blood** and **collect** when there is **no bleeding**. The **tube can be rinsed and reused**.



3. **Use the tubes in order** starting with Tube 1. **Fill the collection tube to the designated level.** To reduce a foamy saliva collection, tap the tube on a hard surface. Insufficient saliva samples cannot be processed.
4. **Replace the cap** tightly to avoid leakage. Record collection start and stop times on the tube label.
5. Place the filled tubes into the biohazard bag and **freeze immediately.** Samples must be frozen a minimum of 2 hours prior to shipping. **Keep sample frozen until ready to ship.**
6. **Repeat these steps for each sample** according to the **Specimen Collection Chart.**



Activate this test online at gdx.net/activate and note the name of the test ordered. Pay close attention to the collection times and amount of saliva required. Failure to do so may cause samples to be rejected or alter results.

SPECIMEN COLLECTION CHART			
Clinicians - Please fill-out the circle for the specimens you want the patient to collect. Only a single result per checkmark is available.			
Tube #1	7AM - 9AM	<input type="radio"/> Estradiol, Salivary <input type="radio"/> Estriol, Salivary* <input type="radio"/> Estrone, Salivary* <input type="radio"/> Testosterone, Salivary <input type="radio"/> Progesterone, Salivary <input type="radio"/> DHEA, Salivary <input type="radio"/> Cortisol	 3ml
Tube #2	11AM - 1PM	<input type="radio"/> Cortisol	 3ml
Tube #3	3PM - 5PM	<input type="radio"/> Cortisol	 3ml
Tube #4	10PM - 12AM	<input type="radio"/> Cortisol	 3ml

STEP 3

Return Collection Pack

1. Confirm each tube has a **completed label attached** with **date of birth, date of collection, collection start time, and collection stop time**. Place the biohazard bag with **tubes** inside the **collection box**.
2. Retain a copy of the **activation number** for future reference **using one of the three labels provided on the bottom of the activation label card**.
3. Visit **gdx.net/activate** to enter the date of your final collection and receive your **confirmation code**. Write the date of collection and your confirmation code on the **activation label card**. Place the **activation card inside** the box.
4. **Close the collection box** and **place** inside the **FedEx shipping bag**. Follow the shipping instructions provided.

